

TRIANGLE-FREE UNIQUELY 3-EDGE COLORABLE CUBIC GRAPHS

SARAH-MARIE BELCASTRO AND RUTH HAAS

ABSTRACT. This paper presents infinitely many new examples of triangle-free uniquely 3-edge colorable cubic graphs. The only such graph previously known was given by Tutte in 1976.

1. HISTORY

Recall that a *cubic* graph is 3-regular, that a *proper 3-edge coloring* assigns colors to edges such that no two incident edges receive the same color, that *edge-Kempe chains* are maximal sequences of edges that alternate between two colors, and that a *Hamilton cycle* includes all vertices of a graph.

It is well known that a cubic graph with a Hamilton cycle is 3-edge colorable, as the Hamilton cycle is even (and thus 2-edge colorable) and its complement is a matching (that can be monochromatically colored). A uniquely 3-edge colorable cubic graph must have exactly three Hamilton cycles, each an edge-Kempe chain in one of the $\binom{3}{2}$ pairs of colors. The converse is not true, as a cubic graph may have some colorings with Hamilton edge-Kempe chains and other colorings with non-Hamilton edge-Kempe chains; examples are given in [12].

The literature classifying uniquely 3-edge colorable cubic graphs is sparse; there is no complete characterization [7]. It is well known that the property of being uniquely 3-edge colorable is invariant under application of $\Delta - Y$ transformations. It was conjectured that every simple planar cubic graph with exactly three Hamilton cycles contains a triangle [13, Cantoni], and also that every simple planar uniquely 3-edge colorable cubic graph contains a triangle [3]. The latter conjecture is proved in [4], where it is also shown that if a simple planar cubic graph has exactly three Hamilton cycles, then it contains a triangle if and only if it is uniquely 3-edge colorable.

Tutte, in a 1976 paper about the average number of Hamilton cycles in a graph [13], offhandedly remarks that one example of a nonplanar triangle-free uniquely 3-edge colorable cubic graph is the generalized Petersen graph $P(9, 2)$, pictured in Figure 1. He describes the graph as two 9-cycles $a_0 \dots a_8$, $b_0 \dots b_8$, with additional edges $a_i b_{2i}$ and index arithmetic done modulo 9.

2010 *Mathematics Subject Classification.* 05C10, 05C15.

Key words and phrases. edge coloring, unique coloring, triangle-free, cubic graphs.

Ruth Haas's work partially supported by NSF grant DMS-1143716 and Simons Foundation Award Number 281291.




FIGURE 1. The generalized Petersen graph $P(9, 2)$ labeled with Tutte's indices.

(The generalized Petersen graph $P(m, 2)$ is defined analogously, and in fact the known cubic graphs with exactly three Hamilton cycles and multiple distinct 3-edge colorings are $P(6k + 3, 2)$ for $k > 1$ [12].) It appears that the search for examples of triangle-free nonplanar uniquely 3-edge colorable cubic graphs ended with Tutte, or at least that any further efforts have been unsuccessful. Multiple sources ([6], [7], [9]) note that Tutte's example is the only known triangle-free nonplanar example. It has been conjectured [3] that $P(9, 2)$ is the only example. In Section 2 we give infinitely many such graphs.

2. NEW EXAMPLES OF TRIANGLE-FREE NONPLANAR UNIQUELY 3-EDGE COLORABLE CUBIC GRAPHS

In [2] the authors introduced the following construction. Consider two cubic graphs G_1, G_2 , and form $G_1 \wr G_2$ by choosing a vertex v_i in G_i ($i = 1, 2$), removing v_i from G_i ($i = 1, 2$), and adding a matching of three edges joining the three neighbors of v_1 with the three neighbors of v_2 . Of course there are many ways to choose v_1, v_2 , and many ways to identify their incident edges, so the construction is not unique. However, it is reversible; given a cubic graph G with a 3-edge cut, we may decompose $G = G_1 \wr G_2$. In that paper we proved the following.

Theorem 2.1. [3.8 of [2]] *Let G_1, G_2 be cubic graphs and a_i the number of 3-edge colorings of G_i . Then $G_1 \wr G_2$ has $a_1 a_2$ edge colorings.*

Define G^\wr to be the infinite family of graphs consisting of all graphs of the form $G \wr G \wr \cdots \wr G$. This leads to the following corollaries of Theorem 2.1.

Theorem 2.2. *If G is a uniquely 3-edge colorable graph, then all graphs in G^\wr are uniquely 3-edge colorable.*




FIGURE 2. A nonplanar, triangle-free, uniquely 3-edge colorable graph with 34 vertices.




FIGURE 3. A nonplanar, triangle-free, uniquely 3-edge colorable graph with 34 vertices that is nonisomorphic to that shown in Figure 2.

Proof. The proof proceeds by induction on the number of copies of G . □

Corollary 2.3. *All members of the infinite family $P(9, 2)^\gamma$ are uniquely 3-edge colorable.*

Note.

In [5], Goldwasser and Zhang proved that if a uniquely 3-edge colorable graph has an edge cut of size 3 or 4 such that each remaining component contains a cycle, then the graph can be decomposed into two smaller uniquely 3-edge colorable graphs. It seems they did not observe the reverse construction.

2.1. Examples and Properties. The smallest member of $P(9, 2)^\gamma$ is of course $P(9, 2)$, which has 18 vertices. For every integer $k > 1$ there are multiple graphs in $P(9, 2)^\gamma$ with $16k + 2$ vertices. Nonisomorphic examples with $k = 2$ are shown in Figures 2 and 3.

The graphs in $P(9, 2)^\gamma$ are clearly all nonplanar. We show next that there are graphs in $P(9, 2)^\gamma$ of every other orientable and nonorientable genus.




FIGURE 4. Embeddings of $P(9, 2)$ on the torus (left) and projective plane (right)

Theorem 2.4. *Every graph in $P(9, 2)^\Upsilon$ with $16k + 2$ vertices has orientable and nonorientable genus at most k . Further, there is a large subfamily of graphs in $P(9, 2)^\Upsilon$, each of which has $16k + 2$ vertices and orientable and nonorientable genus exactly k .*

Proof. We will show by induction that any graph Q_k created using the Υ -construction with k copies of $P(9, 2)$ has orientable and nonorientable genus at most k . The base case holds because $P(9, 2)$ embeds on both the torus (see Figure 4 (left)) and on the projective plane (see Figure 4 (right)).

Now consider Q_k , a graph created using the Υ -construction with k copies of $P(9, 2)$. Q_k was obtained by removing and associating $v \in P(9, 2)$ and some $w \in Q_{k-1}$ via the Υ -construction, where Q_{k-1} is some graph created using $k - 1$ copies of $P(9, 2)$ that has genus $k - 1$ or less by inductive hypothesis. Let \widehat{Q}_k be the graph produced by simply identifying the vertices v and w . The graph \widehat{Q}_k has two blocks that meet at this vertex and by Theorem 1 of [1], the genus of \widehat{Q}_k is the sum of the genera of the blocks (so it is k). Replacing the cut vertex by a 3-edge cut to implement the Υ construction does not increase the genus, which completes the proof of the upper bound on genus.

A copy of a subdivision of $K_{3,3}$ is highlighted in the embedding of $P(9, 2)$ shown in Figure 5. There are four vertices $\{t_1, t_2, t_3, t_4\}$ whose edges are not involved in the subdivided $K_{3,3}$. Any (or all) of $\{t_1, t_2, t_3, t_4\}$ can be removed and the resulting graph will still have a $K_{3,3}$ minor. If Q_k is formed such that in each copy of $P(9, 2)$ only (some) of vertices $\{t_1, t_2, t_3, t_4\}$ are used in the Υ construction then there will still be k disjoint copies of subdivisions of $K_{3,3}$ in Q_k . The genus of a graph is the sum of the genera of its components [1, Cor. 2], so using this construction Q_k has a minor with orientable (resp.




FIGURE 5. $P(9, 2)$ with a copy of a subdivision of $K_{3,3}$ highlighted.

nonorientable) genus exactly k . It is straightforward to draw an embedding of sample Q_k on a surface of orientable or nonorientable genus k .

□

3. CONCLUSION

While we have provided infinitely many examples of triangle-free nonplanar uniquely 3-edge colorable cubic graphs, it is still unknown whether other examples exist. All our examples support Zhang's conjecture [14] that every triangle-free uniquely 3-edge colorable cubic graph contains a Petersen graph minor. That conjecture remains open.

REFERENCES

1. J. Battle, F. Harary, Y. Kodama, and J.W.T. Youngs, *Additivity of the Genus of a Graph*, Bull. Amer. Math. Soc. **68** (1962), 565–568.
2. s.-m. belcastro and R. Haas, *Counting edge-Kempe-equivalence classes for 3-edge-colored cubic graphs*, with R. Haas, Discrete Mathematics, **325** (2014), 77–84.
3. S. Fiorini and R.J. Wilson, *Edge colourings of graphs*, in *Selected topics in graph theory*, ed. L. Beineke and R. Wilson, Academic Press, 1978, 103–126.
4. T.G. Fowler, *Unique Coloring of Planar Graphs*, dissertation, Georgia Institute of Technology, November, 1998; available at people.math.gatech.edu/~thomas/FC/fowlerphd.pdf.
5. J.L. Goldwasser and C.-Q. Zhang, *On minimal counterexamples to a conjecture about unique edge-3-coloring*, Festschrift for C. St. J. A. Nash-Williams, Congr. Numer. **113** (1996), 143–152.
6. J.L. Goldwasser and C.-Q. Zhang, *Uniquely Edge-3-Colorable Graphs and Snarks*, Graphs and Combinatorics **16** (2000), 257–267.
7. J. Gross and J. Yellen, *Handbook of Graph Theory*, CRC Press, Boca Raton, 2004.
8. R. Isaacs, *Infinite families of nontrivial trivalent graphs which are not Tait colorable*, Amer. Math. Monthly **82** (1975), 221–239.
9. T. Jensen and B. Toft, *Graph Coloring Problems*, Wiley Interscience Series, New York, 1995.
10. B. Mohar and C. Thomassen, *Graphs on Surfaces*, Johns Hopkins University Press, Baltimore and London, 2001.
11. N. Robertson, D. Sanders, P. D. Seymour and R. Thomas, *The four colour theorem*, J. Combin. Theory Ser. B. **70** (1997), 2–44.

12. A. Thomason, *Cubic graphs with three hamiltonian cycles are not always uniquely edge colorable*, J. Graph Theory **6** (1982), 219–221.
13. W.T. Tutte, *Hamiltonian circuits*, Colloquio Internazionale sulle Teorie Combinatorics, Atti dei Convegni Lincei. **17** (1976), Accad. Naz. Lincei, Roma I, 193–199.
14. C.-Q. Zhang, *Hamiltonian weights and unique edge-3-colorings of cubic graphs*, J. of Graph Theory **20** (1995), 91–99.

DEPARTMENT OF MATHEMATICS AND STATISTICS, SMITH COLLEGE, NORTHAMPTON,
MA 01063 USA

E-mail address: `smbelcas@toroidalsnark.net`

DEPARTMENT OF MATHEMATICS AND STATISTICS, SMITH COLLEGE, NORTHAMPTON,
MA 01063 USA

E-mail address: `rhaas@smith.edu`